

DRY-CLOX- cloxacillin benzathine gel
Boehringer Ingelheim Vetmedica, Inc

Dry-Clox®
(cloxacillin benzathine)
Intramammary Infusion
FOR USE IN DRY COW ONLY

NADA 055-058, Approved by FDA

Caution:

Federal law restricts this drug to use by or on the order of a licensed veterinarian.

Description:

Dry-Clox (cloxacillin benzathine) for Intramammary Infusion into the Dry Cow is a product which provides bactericidal activity against gram-positive bacteria. The active agent, cloxacillin benzathine, is a sparingly soluble salt of the semisynthetic penicillin, cloxacillin. Cloxacillin is a derivative of 6-aminopenicillanic acid, and therefore is chemically related to other penicillins. It has, however, the antibacterial properties described below, which distinguish it from certain other penicillins.

Each 10 mL disposable syringe contains cloxacillin benzathine equivalent to 500 mg of cloxacillin activity in a stable peanut oil gel. This product was manufactured by a non-sterilizing process.

Storage:

Do not store above 25°C (77°F). Do not freeze.

Action:

In the non-lactating mammary gland, Dry-Clox (cloxacillin benzathine) provides bactericidal levels of the active antibiotic, cloxacillin, for a prolonged period of time. This prolonged activity is due to the low solubility of the cloxacillin benzathine and to the slow-release oil-gel base. This prolonged contact between the antibiotic and the pathogenic organism enhances the probability of a bacteriological cure.

Cloxacillin is not destroyed by the enzyme, penicillinase, and therefore, is active against penicillin-resistant strains of *Staphylococcus aureus*. It is also active against non-penicillinase-producing *Staphylococcus aureus* as well as *Streptococcus agalactiae*.

The class disc, Methicillin 5 mcg, should be used to estimate the *in vitro* susceptibility of bacteria to cloxacillin.

Indications:

For the treatment of mastitis in dairy cows during the dry period.

Dry-Clox has been shown by extensive clinical studies to be efficacious in the treatment of mastitis in dry cows, when caused by *Streptococcus agalactiae* and *Staphylococcus aureus* including penicillin-resistant strains.

Treatment of the dry cow with Dry-Clox is indicated in any cow known to harbor any of these organisms in the udder at drying off, or which has had repeated attacks of mastitis during the previous lactation, or is affected with mastitis at drying off, if caused by susceptible organisms.

Dosage for Dry Cows:

Infuse the contents of one syringe (10 mL) into each quarter following the last milking. See Directions for Use.

Directions for Use:

Dry-Clox (cloxacillin benzathine) is for use in dry cows only. Administer immediately after the last milking. **Use no later than 30 days prior to calving.**

Completely milk out all four quarters. The udder and teats should be thoroughly washed with warm water containing a suitable dairy antiseptic and dried, preferably using individual paper towels. Carefully scrub the teat end and orifice with 70% alcohol, using a separate swab for each teat. **Allow to dry.**

Dry-Clox is packaged with the Opti-Sert[®] Protective Cap.

For partial insertion: Twist off upper portion of the Opti-Sert Protective Cap to expose 3–4 mm of the syringe tip.

For full insertion: Remove protective cap to expose the full length of the syringe tip.

Insert syringe tip into the teat canal and expel the entire contents of syringe into the quarter. Withdraw the syringe and gently massage the quarter to distribute the medication.

Do not infuse contents of the mastitis syringe into the teat canal if the Opti-Sert Protective Cap is broken or damaged.

Precautions:

Because it is a derivative of 6-aminopenicillanic acid, Dry-Clox has the potential for producing allergic reactions. Such reactions are rare; however, should they occur, the subject should be treated with antihistamines or pressor amines, such as epinephrine.

Residue Warnings:

1. **For use in dry cows only.**
2. Not to be used within 30 days of calving.
3. Any animal infused with this product must not be slaughtered for food until 30 days after the latest infusion.

How Supplied:

Dry-Clox (cloxacillin benzathine) Intramammary Infusion is supplied as 10 mL syringes containing 500 mg of cloxacillin activity per syringe. One display carton contains 12 syringes. One pail contains 144 syringes.

NDC 0010-4720-01 - 10 mL syringe

NDC 0010-4720-02 - 12 syringes

NDC 0010-4720-03 - 144 syringes

Opti-Sert is a registered trademark of Zoetis W LLC - used under license.

Dry-Clox is a registered trademark of Boehringer Ingelheim Vetmedica, Inc.

© 2014 Boehringer Ingelheim Vetmedica, Inc. All Rights Reserved.

Made in Italy

Manufactured for:

Boehringer Ingelheim Vetmedica, Inc.

St. Joseph, MO 64506 U.S.A.

51729292

472001-03

Principal Display Panel – Syringe Label

NDC 0010-4720-01

Dry-Clox®

(cloxacillin benzathine)

Intramammary Infusion



Equivalent to 500 mg cloxacillin

Caution: Federal law restricts this drug to use by or on the order of a licensed veterinarian.

FOR USE IN DRY COW ONLY

Net Contents: 10 mL

NADA 055-058, Approved by FDA



NDC 0010-4720-01

Dry-Clox[®]
(cloxacillin benzathine)
**Intramammary
Infusion**

Equivalent to 500 mg cloxacillin
Caution: Federal law restricts this drug
to use by or on the order of a licensed
veterinarian.
FOR USE IN DRY COW ONLY
Net Contents: 10 mL
NADA 055-058, Approved by FDA

Each 10 mL syringe contains 500 mg
cloxacillin activity as cloxacillin benzathine.
This product was manufactured by a
non-sterilizing process.

Dosage: Infuse the contents of one syringe
(10 mL) into each quarter following the last
milking. See package insert for Directions for
Use.


Residue Warnings:

1. For use in dry cows only.
2. Not to be used within 30 days
of calving.
3. Any animal infused with this
product must not be
slaughtered for food until
30 days after the latest
infusion.

Storage: Do not store above 25°C (77°F).
Do not freeze.

Made in Italy 472002-01
51722677

Manufactured for:
Boehringer Ingelheim Vetmedica, Inc.
St. Joseph, MO 64506 U.S.A.



Lot

Exp

Principal Display Panel – 144 syringe pail label

NDC 0010-4720-03

Dry-Clox[®]

(cloxacillin benzathine)

Intramammary Infusion

Equivalent to 500 mg cloxacillin

Caution: Federal law restricts this drug to use by or on the order of a licensed veterinarian.

FOR USE IN DRY COW ONLY

Net Contents: 144 x 10 mL syringes

NADA 055-058, Approved by FDA



NDC 0010-4720-03

Dry-Clox[®]

(cloxacillin benzathine)
Intramammary Infusion

Equivalent to 500 mg cloxacillin

Caution: Federal law restricts this drug to use by or on the order of a licensed veterinarian.

FOR USE IN DRY COW ONLY

Net Contents: 144 x 10 mL syringes

NADA 055-058, Approved by FDA



**Boehringer
Ingelheim**

472009-00
51729192

DRY-CLOX

cloxacillin benzathine gel

Product Information

Product Type	PRESCRIPTION ANIMAL DRUG	Item Code (Source)	NDC:00 10-4720
Route of Administration	INTRAMAMMARY		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
CLOXACILLIN BENZATHINE (UNII: AC79L7PV2G) (CLOXACILLIN - UNII:O6X5QGC2VB)	CLOXACILLIN	500 mg in 10 mL

Inactive Ingredients

Ingredient Name	Strength
PEANUT OIL (UNII: 5TL50QU0W4)	
HYDROGENATED CASTOR OIL (UNII: ZF94AP8MEY)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0010-4720-02	12 in 1 CARTON		
1	NDC:0010-4720-01	10 mL in 1 SYRINGE		
2	NDC:0010-4720-03	144 in 1 PAIL		
2	NDC:0010-4720-01	10 mL in 1 SYRINGE		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NADA	NADA055058	03/11/1975	

Labeler - Boehringer Ingelheim Vetmedica, Inc (007134091)

Revised: 8/2018

Boehringer Ingelheim Vetmedica, Inc