

DORMOSEDAN GEL®

(detomidine hydrochloride)

Prescribed by your veterinarian, DORMOSEDAN GEL® (detomidine hydrochloride) is a safe, effective way for you to administer a mild, standing sedative prior to stressful situations and minor, nonpainful husbandry procedures, including:

- Shoeing
- Body clipping
- Bridle path trimming
- Mane pulling
- Sheath cleaning
- First time turnouts for fractious horses

It's important to withhold food and water until the sedative effect of the product has worn off. When used prior to first time turnouts, supervise horse to prevent grazing until the sedative has worn off.

DORMOSEDAN GEL can only be obtained through a prescription from your horse's veterinarian.

Please follow the dosing guide and instructions below on how to administer DORMOSEDAN GEL.



DOSING REFERENCE GUIDE

The dosing syringe delivers the product in 0.25 mL increments. The following dosing table provides the dose volume to be administered for the corresponding body weight in 0.25 mL increments.

APPROXIMATE BODY WEIGHT (lb)	DOSE VOLUME (mL)
330–439	1.00
440–549	1.25
550–659	1.50
660–769	1.75
770–879	2.00
880–989	2.25
990–1,099	2.50
1,100–1,209	2.75
1,210–1,320	3.00

The duration and level of sedation are dose dependent. The time it takes before your horse is fully sedated is approximately 40 minutes and lasts for about 90 to 180 minutes. If you feel you've given your horse a larger dose than necessary, let him recover in a warm, quiet place and withhold food and water until the sedative has worn off.

Consult your veterinarian for further instructions.

The syringe is designed to be used only once. Partially used syringes should be properly discarded.

HOW TO ADMINISTER DORMOSEDAN GEL



STEP ONE

Put on impermeable gloves and remove the syringe from the outer carton. While holding the plunger, turn the ring-stop on the plunger until the ring is able to slide freely up and down the plunger.




STEP TWO

Position the ring in such a way that the side nearest the barrel is at the desired volume marking. Turn the ring to secure it in place.



STEP THREE

Make sure the horse's mouth contains no feed. Remove the cap from the tip of the syringe and save for cap replacement. Insert the syringe tip into the horse's mouth from the side of the mouth, placing the syringe tip beneath the tongue at the level of the corner of the mouth. Depress the plunger until the ring-stop contacts the barrel, depositing the product under the tongue.



STEP FOUR

Take the syringe out of the horse's mouth, recap the syringe and return it to the outer carton for disposal. Remove and discard gloves or wash them in large amounts of running water.

For best results, allow adequate time (a minimum of 40 minutes) between the administration of DORMOSEDAN GEL and beginning the procedure. In general, horses show sedative effects lasting approximately 90 to 180 minutes.

IMPORTANT SAFETY INFORMATION

DORMOSEDAN GEL is contraindicated in horses with known hypersensitivity to detomidine. Intravenous potentiated sulfonamides should not be used in anesthetized or sedated horses, as potentially fatal dysrhythmias may occur. Do not use DORMOSEDAN GEL in horses with pre-existing atrioventricular (AV) or sinoatrial (SA) blocks, cardiovascular disease, respiratory disorders, liver or kidney diseases, or in conditions of shock, severe debilitation or stress due to extreme heat, cold, fatigue or high altitude. Appropriate precautions should be taken while handling and using gel-dosing syringes, as DORMOSEDAN GEL can be absorbed following direct exposure to skin, eyes or mouth, and may cause irritation. The use of impermeable gloves is advised. Please see the full prescribing information, or go to www.DormosedanGel.com.



