Lactated Ringer's



Injection

This sterile, nonpyrogenic solution is indicated for the correction of electrolyte depletion, metabolic acidosis, and dehydration of cattle, calves, horses, sheep and swine. May be injected intravenously, subcutaneously or intraperitoneally (except in horses) using strict aseptic technique. Packaged in a hard plastic bottle that has a fold-flat integrated hook for hanging. Prescription.

ITEM#	DESCRIPTION	SIZE
501033	Lactated Ringer's Injection	Liter







VetOne products are sold exclusively through MWI Animal Health sea 800.824.3703 mwiah.com For product information, log on to your MWI website account or visit vetone.net

Lactated Ringer's



VET ONE[®] Lactated Ringer's

Injection

Sterile - Nonpyrogenic solution

Caution: Federal law restricts this drug to use by or on the order of a licensed veterinarian.

Warning: For animal use only.

Keep out of reach of children.

Solution should be warmed to body temperature prior to administration and administered at a slow rate. This is a single dose unit. It contains no preservatives. If entire contents are not used, discard unused portion.

Indications: For the correction of electrolyte depletion, metabolic acidosis and dehydration of cattle, calves, horses, sheep, and swine.

Dosage and Administration: May be injected intravenously, subcutaneously or intraperitoneally (except in horses) using strict aseptic technique.

Cattle and Horses: 2 to 5 mL per pound of body weight, depending on size and condition of animal, repeated 1 to 3 times daily or as needed.

Swine and Sheep: 2 to 5 mL per pound of body weight depending on size and condition of animal, repeated 1 to 3 times as needed.

If administered subcutaneously divide dosage in several sites of injection and massage points of injection to aid in absorption and help prevent inflammation and /or sloughing.

Each 100 mL contains:

Sodium Chloride 600 mg
Sodium Lactate 310 mg
Potassium Chloride 30 mg
Calcium Chloride Dihydrate
Water or Injection q.s

Warning: Do not administer to horses by intraperitoneal injection. Do not administer to animals with inadequate renal function. Not for us in lactic acidosis.

Storage Information: Store between 15° and 30°C (59°F-86°F).

Rev 12/17



VetOne products are sold exclusively through MWI Animal Health Section 800.824.3703 mwiah.com For product information, log on to your MWI website account or visit wetone.net

